






Answers to Explore the Math

- A.** The polynomial expressions all involve only the sum of constant multiples of non-negative integer powers of x .
- B.** Some have negative or fractional powers of x . Others have an x in the denominator of a fraction. Still others have x 's and y 's, and one even involves a sine function.
- C.** Answers may vary. A polynomial expression is an expression that is the sum of constant multiples of non-negative integer powers of x .

D.

Polynomial Function	Type	Sketch of Graph	Description of Graph	Domain and Range	Existence of Asymptotes?
$f(x) = x$	linear		The graph is a line through (0, 0) at a 45° angle to the x-axis.	$D = \{x \in \mathbf{R}\}$ $R = \{f(x) \in \mathbf{R}\}$	None
$f(x) = x^2$	quadratic		The graph is a parabola through (0, 0) opening up.	$D = \{x \in \mathbf{R}\}$ $R = \{f(x) \in \mathbf{R} \mid f(x) \geq 0\}$	None
$f(x) = x^3$	cubic		The graph increases from $-\infty$ to ∞ and flattens out around the origin.	$D = \{x \in \mathbf{R}\}$ $R = \{f(x) \in \mathbf{R}\}$	None

$f(x) = x^4$	quartic		The graph looks like the graph of x^2 only much flatter near the origin.	$D = \{x \in \mathbf{R}\}$ $R = \{f(x) \in \mathbf{R} \mid f(x) \geq 0\}$	None
$f(x) = x^5$	quintic		The graph looks like the graph of x^3 only much flatter near the origin.	$D = \{x \in \mathbf{R}\}$ $R = \{f(x) \in \mathbf{R}\}$	None

E. The linear, cubic, and quintic behave similarly and all have odd powers, while the quadratic and quartic behave similarly and have even powers.

F.

$f(x) = x$	Δ_1
$f(-3) = -3$	
$f(-2) = -2$	1
$f(-1) = -1$	1
$f(0) = 0$	1
$f(1) = 1$	1
$f(2) = 2$	1
$f(3) = 3$	1

$f(x) = x^2$	Δ_1	Δ_2
$f(-3) = 9$		
$f(-2) = 4$	-5	2
$f(-1) = 1$	-3	2
$f(0) = 0$	-1	2
$f(1) = 1$	1	2
$f(2) = 4$	3	2
$f(3) = 9$	5	

$f(x) = x^3$	Δ_1	Δ_2	Δ_3
$f(-3) = -27$			
$f(-2) = -8$	19	-12	
$f(-1) = -1$	7	-6	6
$f(0) = 0$	1	0	6
$f(1) = 1$	1	6	6
$f(2) = 8$	7	12	6
$f(3) = 27$	19		

$f(x) = x^4$	Δ_1	Δ_2	Δ_3	Δ_4
$f(-3) = 81$				
$f(-2) = 16$	-65			
$f(-1) = 1$	-15	50	-36	
$f(0) = 0$	-1	14	-12	24
$f(1) = 1$	1	2	12	24
$f(2) = 16$	15	14	36	24
$f(3) = 81$	65	50		

$f(x) = x^5$	Δ_1	Δ_2	Δ_3	Δ_4	Δ_5
$f(-3) = -243$					
	211				
$f(-2) = -32$		-180			
	31		150		
$f(-1) = -1$		-30		-120	
	1		30		120
$f(0) = 0$		2		0	
	1		30		120
$f(1) = 1$		30		120	
	31		150		
$f(2) = 32$		180			
	211				
$f(3) = 243$					

The number of finite differences to get down to a constant is determined by the degree of the polynomial.

Answers to Reflecting

- K. Both are sums of constant multiples of non-negative integer powers of x .
- L. As the degree increases, the graph flattens out near the origin and becomes steeper away from the origin. The number of finite differences needed to obtain a constant increases with the degree.